



UNIVERSITY OF
BIRMINGHAM

SCHOOL OF
PHYSICS AND
ASTRONOMY

University of Birmingham new simulation framework update

P.P. Allport, L. Gonella, P. Ilten, P.G. Jones, P.R. Newman,
H. Wennlöf


7th of May 2020

h.wennlof@cern.ch




ESCalate framework (G4E+eJANA)

- Current working status:
 - G4E+eJANA (ESCalate) set up locally.
 - Can change geometry and active detector parts in G4E.
 - SVT implemented analogously to previous EICROOT studies (<http://cern.ch/go/xKk6>).
 - Stave construction set up based on EICROOT staves in a parameterised way, where it is easy to vary material thicknesses. Material scans made to create three stave types; inner ($0.3\% X_0$), outer ($0.8\% X_0$), and time-stamping ($1.6\% X_0$). Staves made to level of "simple structure".




- Staves consist of
 - Kapton
 - Aluminium
 - Silicon (active volume)
 - Carbon fiber
 - Water cooling pipes




ESCalate framework (G4E+eJANA)

- Current working status:
 - Particle gun constructed, generating same distributions as used in previous EICROOT simulations.
 - Can write analysis code using eJANA plugin system.
 - Plugin written to create ROOT tree with the same structure as was done in EICROOT, so same final analysis tools can be used (making comparison straightforward).
 - Have set up Pythia generator. Can generate events, and propagate them through G4E geometry. Hit positions can be extracted.
- Vertex reconstruction currently **not** working
 - ACTS implementation underway, will reconstruct both tracks and vertices. ESCalate developers working with ACTS developers to get it functioning




Preliminary results – GenFit momentum reconstruction vs EICROOT

- Parameters used:
 - Particle: π^+
 - Transverse momentum range: 0 to 5 GeV/c
 - Pseudorapidity range: $-0.5 \leq \eta \leq 0.5$
 - Default pixel size: $20 \times 20 \mu\text{m}^2$
 - Magnetic field: uniform 1.5 T
- Relative momentum resolution can be extracted (at innermost layer)
 - G4E/GenFit values correct order of magnitude.
 - No TPC/extra silicon layers present, which degrades the resolution compared to previously presented studies.



Relative momentum resolution vs momentum



Relative momentum resolution vs η

Planned work - ESCalate

- Benchmarking against EICROOT results once vertex reconstruction is in place.
 - Analysis chain set up
 - Once reconstruction works, everything is ready to get results for Pavia workshop
- Physics simulations, once basic simulations yield reliable and consistent results.

Planned work - EICROOT

- Simulate best layouts with
 - New beampipe configuration (i.e. 31 mm radius)
 - 1.5 T and 3 T magnetic field comparison in barrel region
 - More realistic TPC, as simulated by eRD6
 - If approximate EIC parameters are decided on, please let me know; h.wennlof@cern.ch
- Study impact of adding a third inner layer, to be able to reconstruct even if a layer is missed
- Study impact of thinner inner layers, more ITS3-like

